

523
NPSOR-91-11

NAVAL POSTGRADUATE SCHOOL

Monterey, California



SERVICE-ADAPTIVE MULTI-TYPE REPAIRMAN PROBLEMS

Donald P. Gaver
John A. Morrison
Rogerio Silveira

March 1991

Approved for public release; distribution is unlimited.

Prepared for:

Defense Communications Agency, Reston, Virginia, and
ONR, Arlington, VA (Direct Funding at NPS)

Fedders
D 208.14/2
NPS-OR-91-11

NAVAL POSTGRADUATE SCHOOL,
MONTEREY, CALIFORNIA

Rear Admiral R. W. West, Jr.
Superintendent

Harrison Shull
Provost

This report was prepared for Defense Communications Agency, Reston, Virginia, and
ONR, Arlington, VA (Direct Funding at NPS).

This report was prepared by:

Unclassified

Security Classification of this page

REPORT DOCUMENTATION PAGE

1a Report Security Classification UNCLASSIFIED		1b Restrictive Markings	
2a Security Classification Authority		3 Distribution Availability of Report	
2b Declassification/Downgrading Schedule		Approved for public release; distribution is unlimited	
4 Performing Organization Report Number(s) NPSOR-91-11		5 Monitoring Organization Report Number(s)	
6a Name of Performing Organization Naval Postgraduate School	6b Office Symbol (If Applicable) OR	7a Name of Monitoring Organization Defense Communications Agency, Reston, Virginia and ONR, Arlington, VA	
6c Address (city, state, and ZIP code) Monterey, CA 93943-5000		7b Address (city, state, and ZIP code)	
8a Name of Funding/Sponsoring Organization Naval Postgraduate School	8b Office Symbol (If Applicable)	9 Procurement Instrument Identification Number O&MN, Direct Funding	
8c Address (city, state, and ZIP code) Monterey, California		10 Source of Funding Numbers	
		Program Element Number	Project No
		Task No	Work Unit Accession No
11 Title (Include Security Classification) Service-Adaptive Multi-Type Repairman Problems			
12 Personal Author(s) Gaver, D. P., Morrison, J. A., and Silveira, R			
13a Type of Report Technical	13b Time Covered From To	14 Date of Report (year, month, day) 1991, March	15 Page Count
16 Supplementary Notation The views expressed in this paper are those of the author and do not reflect the official policy or position of the Department of Defense or the U.S. Government.			
17 Cosati Codes		18 Subject Terms (continue on reverse if necessary and identify by block number)	
Field	Group	Subgroup	
19 Abstract (continue on reverse if necessary and identify by block number)			
<p>The classical "repairman problem," cf. Feller (1967) is generalized to consider r failure-prone machine types, each type having its own individual failure rate and also repair rate. Each failed machine joins its type queue, and is repaired by a single server. Several dynamic service priority schemes are considered that approximate first-come first-served, longest-line first, and least-available first. A heavy-traffic asymptotic analysis determines approximations to the time-dependent mean and covariance of individual type queue lengths, and shows that the marginal joint distribution of queue lengths is approximately Ornstein-Uhlenbeck. Numerical illustrations of approximation accuracy are provided, as are suggested applications to computer performance and manufacturing systems analysis.</p>			
20 Distribution/Availability of Abstract		21 Abstract Security Classification	
<input checked="" type="checkbox"/> unclassified/unlimited <input type="checkbox"/> same as report <input type="checkbox"/> DTIC users		Unclassified	
22a Name of Responsible Individual D. P. Gaver		22b Telephone (Include Area code) (408) 646-2605	22c Office Symbol OR/Gv

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted

All other editions are obsolete

security classification of this page

Unclassified

SERVICE-ADAPTIVE MULTI-TYPE REPAIRMAN PROBLEMS

Donald P. Gaver,
Naval Postgraduate School,
Monterey, CA 93943

John A. Morrison,
AT&T Bell Laboratories,
Murray Hill, NJ 07974

Rogério Silveira
Centre de Análises de Sistemas Navais (CASNAV)
Avenida Pasteur 480, Fdas.
Rio de Janeiro, RJ 22290, Brazil

Abstract. The classical "repairman problem," cf. Feller (1967) is generalized to consider r failure-prone machine types, each type having its own individual failure rate and also repair rate. Each failed machine joins its type queue, and is repaired by a single server. Several dynamic service priority schemes are considered that approximate first-come first-served, longest-line first, and least-available first. A heavy-traffic asymptotic analysis determines approximations to the time-dependent mean and covariance of individual type queue lengths, and shows that the marginal joint distribution of queue lengths is approximately Ornstein-Uhlenbeck. Numerical illustrations of approximation accuracy are provided, as are suggested applications to computer performance and manufacturing systems analysis.

1. INTRODUCTION

The classical “repairman problem” described by Feller (1967), Cox and Smith (1961), and in all later texts on applied probability, is a well-motivated example of a finite Markov chain in continuous time. Not only is it a plausible first model for the productivity of a group of similar machines that are subject to failure and susceptible to repair by one (or more) repairmen, as described by Feller and subsequent authors, but it provides initial rough models for prototypical computer science situations such as multiprocessing and polling. The purpose of this paper is to extend the basic simple *single-type* (of machine, for instance) Markov repairman model to various *multi-type* models, i.e., ones in which several types of items (e.g., machines or computer system users) become applicants for service. These then *may* join a single queue and be serviced in arrival order: first-in, first-out or FIFO. Alternatively, the service applicant types can be individually queued, and a service assignment can be made from one of these queues at a suitable instant, such as at the end of a service; evaluation of the latter *dynamic complete-service assignment* rules are emphasized in this paper. Note parenthetically that in principle there need be no restriction to choice from among the waiting types *only* when service is completed: preemption at some intermediate time can be an effective tactic, particularly if setup or switchover times or costs are not excessive, in order to focus attention on a machine type that momentarily has a long queue and thus is in excessively short supply. One limit of preemption is some form of processor sharing, in which the server’s tendency to service a particular type among service applicants depends moment by moment upon lengths of queues, or urgency. In some

contexts switching back and forth between partially completed services is severely penalized by substantial switchover times, which reduces the attractiveness of the tactic. It is even possible that such switching can lead to bistability, an elementary form of chaos; see Jaiswal, *et al.* (1990). We do not consider such switching models in the present paper.

In what follows we first describe our mathematical model, paying particular attention to a handy probabilistic device for representing purposeful and adaptive selection of a new service incumbent just after the previous one completes service and leaves. An important example could be the simple practice of selecting from the longest queue; another is selecting from the queue of items that is least available. We then introduce a state space scaling that is appropriate when the number of each machine type grows large and the system is in heavy traffic, so the server is essentially never idle. Asymptotic expansions are introduced in terms of a large parameter, a ; a is of the order of magnitude of the number of machines. These expansions reveal the approximate time-dependent means of the lengths of the queues awaiting processing, and also the time-dependent joint characteristic function (ch.f.) of the random noise terms that characterize departures from the mean. The latter approaches the ch.f. of a multivariate Ornstein-Uhlenbeck process; see Arnold (1974). The scaling studied is the same as that of McNeil and Schach (1973) but the present results pertain specifically to a multi-type queueing system's transient behavior. Finally, numerical results computed from the model by numerical solutions of systems of differential equations are compared to simulations, with good agreement exhibited. The computational time for the presently described

analytical-numerical procedure is a small fraction of that required to conduct a comparable simple simulation, suggesting its practicality and usefulness when appropriate.

2. THE MODEL

In this section we describe the initial basic model for a multi-type repairman problem with adaptive service.

a) Demands

There are K_j items ("machines") of type j , each of which, when operating, fail independently at Markovian rate λ_j ; $j = 1, 2, \dots, r$. In fact, the failure rate can even be smoothly time-dependent, with failure rate at time (*not* age) t equal to $\lambda_j(t)$, but we emphasize discussion of the time-homogeneous situation here.

b) Service

There is a single server. When an item of type j is undergoing service its repair process is Markovian with rate ν_j . Again, time-dependence is allowable. We only consider the case of *heavy traffic*, which means that $\sum_{j=1}^r K_j \lambda_j / \nu_j > 1$ when rates are constant. This assumption assures that queues for all items form soon after the process begins; the queues grow towards steady-state levels and fluctuate but essentially never dissipate.

c) Service Discipline; Models for Adaptation

Service is organized by assembling different demand types in their own queues. Let $N_j(t)$ denote the number of items of type j awaiting service at time t ; if a type j item happens to be undergoing service it is also included in the count for $N_j(t)$. Let $I(t)$ be an indicator variable; $I(t) = i$ ($i = 1, 2, \dots, r$) means that an item of type i is in service at time t .

When an item of any type completes service its replacement is *modeled* as if selected by a random device: given that service is completed at time t and the number of items in the various queues is $\underline{N}(t) = \underline{n}$ then with probability $q_j(\underline{n}, \underline{K}, \underline{w}(t))$ the new server incumbent at t is of type j . The function $q_j(\cdot)$ is to be chosen so as to model desired service protocols; it is, in particular, assumed to be homogeneous of degree zero in \underline{n} and \underline{K} . We select for study specific forms that represent, for example, (i) first-in, first-out (FIFO) service selection, (ii) longest-line-next (LOLIN), (iii) least-available-next (LAIN) or any weighted combination thereof. Here are the explicit forms chosen; for $j = 1, 2, \dots, r$

$$(i) \text{ FIFO: } q_j(\underline{n}(t), \underline{K}, \underline{w}(t)) = \frac{n_j(t)w_j(t)}{\sum_k n_k(t)w_k(t)} \quad (2.1)$$

$$(ii) \text{ LOLIN: } q_j(\underline{n}(t), \underline{K}, \underline{w}(t)) = \frac{(n_j(t))^p w_j(t)}{\sum_k (n_k(t))^p w_k(t)}, p \gg 1 \quad (2.2)$$

$$(iii) \text{ LAIN: } q_j(\underline{n}(t), \underline{K}, \underline{w}(t)) = \frac{(K_j - n_j(t))^{-p} w_j(t)}{\sum_k (K_k - n_k(t))^{-p} w_k(t)}, p \gg 1 \quad (2.3)$$

where $(w_j(t)) = \underline{w}(t)$ is a set of positive weights and p is a positive tuning constant (suitably chosen to be "large, but not too large"). Note that each specific function is the ratio of homogeneous functions, differentiable in n_j and also K_j . To motivate (2.3), it is clear that if, say, the availability of item 1, namely $K_1 - n_1$ is less than that of any other item $j \neq 1$, then its *anti-availability* $(K_1 - n_1)^{-1}$ is greater than that of any other, and the difference is enhanced by

increasing p in $(K_1 - n_1)^{-p}$; consequently the q_j of (2.3) increases controllably the probability of choosing item j for next service if it is least available. In principle ambiguity exists if $n_k(t) = K_k$ for more than one machine; we do not consider this unlikely event. An analogous argument holds for (2.2). It has long been understood that (2.1) with equal weights already well-approximates FIFO; its use eliminates the need for a far more elaborate state space to describe true FIFO. See Gaver and Morrison (1991) for a treatment of our FIFO model by somewhat different methods.

Although we model the selection of the next item to be served by a random process governed by q_j the intention is that this model emulate deterministic selection. Modeling by a probability that is a sufficiently smooth (differentiable) function of state variables $n_j(t)$ permits computation of asymptotic expansions that conveniently characterize measures of the probabilistic fluctuations of those state variables; in particular the approximate variance-covariance function of $\underline{N}(t)$. It will be shown by simulation for specific examples that numerical agreement of our probabilistic analytical models is very close to the corresponding deterministic selection results. The computing time for obtaining the numerical results from the analytical model is, however, a small fraction of the time to obtain the simulation results. This allows considerable elaboration of the present basic model; for some details see Gaver *et al.* (1991).

There follows a discussion of application areas for our class of models.

Adaptive Polling

Suppose a single processor, e.g., computer or database, receives tasks from r different classes of input devices which we will call terminals. When a

terminal in class j makes a demand it remains quiescent until that demand is served, after which it becomes active again at rate λ_j . The requests from each terminal individually queue for service at the processor; when a request or task is accepted by the (single) processor it is served to completion; the server is said to **poll** terminal class queues for the next request or task to be served when completion occurs. The analysis herein permits assessment of various polling selection policies, such as one that dynamically favors the longest queue (equally-weighted LOLIN) and thus tends to equalize queue lengths and waits for service. If the terminal classes were to be information sources in a military command-control-intelligence (C²I) system, and requests were database updates, the priority weights, w_j , could be adjusted to account for the current credibility of inputs by classes of terminals; the most credible would be processed first in times of stress and heavy usage, as during combat. It is easily possible to adjust the service policy to that of class-preferential processor sharing, see Gaver and Jacobs (1986) and Morrison (1987) for similar asymptotic studies.

Why LAIN? Promotion of Item-Cooperative Service in Reliability or Manufacturing Applications

Several real situations call for the use of the least-available-next (LAIN) policy as at least attractive, if possibly not demonstrably optimal. Consider the modeling of adaptive service to achieve

System Availability

First, suppose that exactly one of each item type is needed in order to render fully operational a *platform system* such as a fully-equipped aircraft. Then the least-available item type controls the number of platforms available for action. In case there are fewer of each item type than there are platforms then all available platforms will be in action at any time and the simple process described above is in play and its analysis is appropriate. For a more detailed description of the aircraft availability process see Gaver *et al.* (1991).

Manufacturing: Stage-Wise Production

Second, suppose instead that each item type represents a machine that is a stage in some sequential manufacturing process. Manufacturing, or other processing, proceeds in successive stages, starting at a bank of (available) machines of type 1, the output of that stage proceeding to, and through, machines of type 2, *etc.*, through machines of type r at Stage r . Machines are, however, susceptible to failure and only those that have not failed (are available) are productive. Provided that all machine banks are about equally productive, so that there are no production bottlenecks in the absence of failures, it is of interest to schedule repairs to maintain that balance when realistic failures do occur. Certainly the LAIN policy has the desired effect of

maintaining equality among machine types available. A generalization might well be to induct for next service that machine type that has the largest queue of in-process production items awaiting that particular machine bank.

3. MARKOV STRUCTURE AND ASYMPTOTIC ANALYSIS

It is clear from the assumptions made that the process $\{\underline{N}(t), I(t)\} = \{N_1(t), N_2(t), \dots, N_r(t), I(t)\}$ is Markov in continuous time with a finite discrete state space. If all parameters are actually independent of time such a process will ultimately reach steady-state, exhibiting a stationary distribution. But explicit calculation of that distribution, or of the distribution at any finite time, t , is an involved exercise. Special efforts would be required for system design and operation. The same is true if simulation is used to study the behavior of various scheduling options such as FIFO, LOLIN, or LAIN. Hence the approach of approximating the behavior of the above multivariate process by the sum of a deterministic mean and a continuous-state-space diffusion process with Gaussian marginal distributions is attractive. McNeil and Schach (1973) carried out a comparable program for a wide variety of models, but our approach and results differ from theirs in various respects; in particular our process is multivariate.

Asymptotic Analysis

Let

$$P_\ell(\underline{n}, t) = P\{N_1(t) = n_1, \dots, N_r(t) = n_r, I(t) = \ell\} \equiv P\{\underline{N}(t) = \underline{n}, I(t) = \ell\}, \quad (3.1)$$

where initial conditions have been suppressed. Then standard arguments lead to the forward Kolmogorov equations

$$\begin{aligned} \frac{\partial}{\partial t} P_\ell(\underline{n}, t) + \sum_{j=1}^r \lambda_j (K_j - n_j) P_\ell(\underline{n}, t) + v_\ell P_\ell(\underline{n}, t) = \\ \sum_{j=1}^r \lambda_j (K_j - n_j + 1) P_\ell(\underline{n} - \underline{e}_j, t) + q_\ell(\underline{n}, K) \sum_{j=1}^r v_j P_j(\underline{n} + \underline{e}_j, t), \end{aligned} \quad (3.2)$$

for $n_\ell \neq 0$, $\underline{n} \neq \underline{e}_\ell$ and $0 \leq \underline{n} \leq \underline{k}$, where $\underline{e}_j = (0, 0, \dots, 1, 0, 0)$, i.e., 1 occurs only in the j^{th} entry of vector \underline{e}_j . We do not include the special equations appropriate when $\underline{n} = \underline{0}$ and $\underline{n} = \underline{e}_\ell$ since the heavy traffic assumption renders this unnecessary.

Scaling

For a large value a , i.e., $a \gg 1$ we assume $K_j = a\alpha_j$, $v_j = a\mu_j$. Consider the transformed variables

$$V_j(t) = (N_j(t) - a\beta_j(t)) / \sqrt{a}, j = 1, 2, \dots, r \quad (3.3)$$

wherein $\beta_j(t)$ are deterministic functions to be determined; $a\beta_j(t) \approx E[N_j(t)]$. Define the characteristic function (ch.f.) of $V_j(t)$ for given a as

$$\begin{aligned} \omega_\ell(y, t; a) &= E\left[\exp(iy\underline{V}(t)); I(t) = \ell\right] \\ &= \sum_{(\underline{n})} \exp\left[iy(\underline{n} - a\underline{\beta}(t)) / \sqrt{a}\right] P_\ell(\underline{n}, t). \end{aligned} \quad (3.4)$$

The objective will be to derive asymptotic expansions in powers of $1/\sqrt{a}$ that characterize the mean function $\underline{\beta}(t) = (\beta_1(t), \beta_2(t), \dots, \beta_r(t))$ and also $\underline{\omega} = (\omega_1, \omega_2, \dots, \omega_r)$; from the latter variances and covariances of queue lengths can be deduced. From the limiting characteristic function it may be deduced that the process $\underline{V}(t)$ is approximately multivariate Ornstein-Uhlenbeck, with parameters specific to the particular scheduling rule in force.

Preliminaries

According to its definition, (3.4),

$$\omega_\ell(y, t; a) = \sum_{(\underline{n})} e^{iy(\underline{n} - a\underline{\beta}(t)) / \sqrt{a}} P_\ell(\underline{n}, t) \equiv \sum_{(\underline{n})} e^{iy\underline{v}(\underline{n}, t)} P_\ell(\underline{n}, t) \quad (3.5)$$

where

$$\underline{v}(\underline{n}, t) = (\underline{n} - a\underline{\beta}(t)) / \sqrt{a}. \quad (3.6)$$

Now multiply (3.2) by $e^{iy\underline{v}}$ and sum on \underline{n} . Note that differentiation of (3.5) gives

$$\frac{\partial \omega_\ell}{\partial t} = -i\sqrt{a} \underline{y} \frac{d\underline{\beta}}{dt} \omega_\ell + \sum_{(\underline{n})} e^{iy\underline{v}} \frac{\partial}{\partial t} P_\ell(\underline{n}, t), \quad (3.7)$$

and that the scaling (3.6) implies

$$\begin{aligned} \sum_{(\underline{n})} e^{iy\underline{v}} \sum_{j=1}^r n_j P_\ell(\underline{n}, t) &= \sum_{j=1}^r \sum_{(\underline{n})} (\sqrt{a} v_j + a\beta_j(t)) e^{iy\underline{v}} P_\ell(\underline{n}, t) \\ &= \sum_{j=1}^r \left(-i\sqrt{a} \frac{\partial \omega_\ell}{\partial y_j} + a\beta_j(t) \omega_\ell \right), \end{aligned} \quad (3.8)$$

so from the Kolmogorov equation

$$\begin{aligned} \frac{\partial \omega_\ell}{\partial t} + i\sqrt{a} \underline{y} \frac{d\underline{\beta}}{dt} \omega_\ell + \sum_{j=1}^r \lambda_j \left[K_j \omega_\ell + i\sqrt{a} \frac{\partial \omega_\ell}{\partial y_j} - a\beta_j(t) \omega_\ell \right] + a\mu_\ell \omega_\ell = \\ \sum_{j=1}^r \lambda_j \left[K_j \omega_\ell + i\sqrt{a} \frac{\partial \omega_\ell}{\partial y_j} - a\beta_j \omega_\ell \right] e^{iy_j/\sqrt{a}} + \sum_{(\underline{n})} e^{iy\underline{v}} q_\ell(\underline{n}, \underline{K}, \underline{w}) \sum_{j=1}^r v_j P_j(\underline{n} + \underline{e}_j, t). \end{aligned} \quad (3.9)$$

Expand q_j by noting that in representation (2.1)–(2.3) numerator and denominator are homogeneous functions, so cancellation of \sqrt{a} provides

$$q_j(\underline{n}, \underline{K}, \underline{w}(t)) = q_j(\underline{\beta}(t), \underline{\alpha}, \underline{w}(t)) + \frac{1}{\sqrt{a}} \sum_{k=1}^r v_k \frac{\partial q_j}{\partial \beta_k} + O(1/a). \quad (3.10)$$

Now insert (3.10) into the final sum of (3.9) to get

$$\begin{aligned}
& \frac{\partial \omega_\ell}{\partial t} + i\sqrt{a}\omega_\ell \underline{y} \frac{d\beta}{dt} + \sum_{j=1}^r \lambda_j \left[a\alpha_j \omega_\ell + i\sqrt{a} \frac{\partial \omega_\ell}{\partial y_j} - a\beta_j(t)\omega_\ell \right] + a\mu_l \omega_l = \\
& \sum_{j=1}^r \lambda_j \left[a\alpha_j \omega_\ell + i\sqrt{a} \frac{\partial \omega_\ell}{\partial y_j} - a\beta_j(t)\omega_\ell \right] e^{iy_j/\sqrt{a}} \\
& + aq_\ell(\underline{\beta}, \underline{\alpha}, \underline{w}) \cdot \sum_{j=1}^r \mu_j \omega_j e^{-iy_j/\sqrt{a}} - i\sqrt{a} \sum_{j=1}^r \mu_j \sum_{k=1}^r \frac{\partial q_\ell}{\partial \beta_k} e^{-iy_j/\sqrt{a}} \frac{\partial \omega_j}{\partial y_k} + O(1), \quad (3.11)
\end{aligned}$$

omitting terms of order $1/a$ in the expansion of q_j . Next divide through (3.11) by a and note that

$$\begin{aligned}
& \mu_\ell \omega_\ell - q_\ell \sum_{j=1}^r \mu_j \omega_j e^{-iy_j/\sqrt{a}} + \sum_{j=1}^r \left[\lambda_j (\alpha_j - \beta_j) \left(1 - e^{iy_j/\sqrt{a}} \right) + \frac{i}{\sqrt{a}} \frac{d\beta_j}{dt} y_j \right] \omega_\ell \\
& + \frac{i}{\sqrt{a}} \sum_{j=1}^r \mu_j \sum_{k=1}^r \frac{\partial q_\ell}{\partial \beta_k} \frac{\partial \omega_j}{\partial y_k} e^{-iy_j/\sqrt{a}} + \frac{i}{\sqrt{a}} \sum_{j=1}^r \lambda_j \frac{\partial \omega_\ell}{\partial y_j} \left(1 - e^{iy_j/\sqrt{a}} \right) = O(1/a). \quad (3.12)
\end{aligned}$$

Expand the exponentials to find

$$\begin{aligned}
& \mu_\ell \omega_\ell - q_\ell \sum_{j=1}^r \mu_j \omega_j + \frac{i}{\sqrt{a}} \sum_{j=1}^r \left[\frac{d\beta_j}{dt} - \lambda_j (\alpha_j - \beta_j) \right] y_j \omega_\ell = \\
& - \frac{i}{\sqrt{a}} \sum_{j=1}^r \mu_j \left[q_\ell \omega_j y_j + \sum_{k=1}^r \frac{\partial q_\ell}{\partial \beta_k} \frac{\partial \omega_j}{\partial y_k} \right] + O(1/a). \quad (3.13)
\end{aligned}$$

If (3.11) is summed on ℓ then, noting that since $\sum_{l=1}^r q_l(\underline{\beta}, \underline{\alpha}) = 1$, so $\sum_{\ell=1}^r \partial q_\ell / \partial \beta_k = 0$, etc., there results

$$\begin{aligned}
& \sum_{j=1}^r \mu_j \left(1 - e^{-iy_j/\sqrt{a}} \right) \omega_j + \sum_{j=1}^r \left[\lambda_j (\alpha_j - \beta_j) \left(1 - e^{iy_j/\sqrt{a}} \right) + \frac{i}{\sqrt{a}} \frac{d\beta_j}{dt} y_j \right] \sum_{l=1}^r \omega_l \\
& + \frac{i}{\sqrt{a}} \sum_{j=1}^r \lambda_j \left(1 - e^{iy_j/\sqrt{a}} \right) \sum_{\ell=1}^r \frac{\partial \omega_\ell}{\partial y_j} + \frac{1}{a} \sum_{\ell=1}^r \frac{\partial \omega_\ell}{\partial t} = 0. \quad (3.14)
\end{aligned}$$

The fact that this summed equation holds follows directly from the summed Kolmogorov equation; details are omitted. Now expand the exponentials in powers of $1/\sqrt{a}$ to find

$$\begin{aligned} & \frac{i}{\sqrt{a}} \left\{ \sum_{j=1}^r \mu_j \omega_j y_j + \sum_{j=1}^r \left[\frac{d\beta_j}{dt} - \lambda_j (\alpha_j - \beta_j) \right] y_j \left(\sum_{\ell=1}^r \omega_\ell \right) \right\} \\ & + \frac{1}{a} \left\{ \sum_{j=1}^r \mu_j \omega_j \frac{y_j^2}{2} + \sum_{j=1}^r \lambda_j (\alpha_j - \beta_j) \frac{y_j^2}{2} \left(\sum_{\ell=1}^r \omega_\ell \right) + \sum_{j=1}^r \lambda_j y_j \left(\sum_{\ell=1}^r \frac{\partial \omega_\ell}{\partial y_j} \right) + \sum_{\ell=1}^r \frac{\partial \omega_\ell}{\partial t} \right\} \\ & = O(1/a^{3/2}). \end{aligned} \quad (3.15)$$

Expansions

Next introduce the asymptotic expansion

$$\omega_\ell(\underline{y}, t; a) = \omega_\ell^{(0)}(\underline{y}, t) + \frac{1}{\sqrt{a}} \omega_\ell^{(1)}(\underline{y}, t) + \frac{1}{a} \omega_\ell^{(2)}(\underline{y}, t) + \dots \quad (3.16)$$

Insert (3.16) into (3.13) and isolate terms of $O(1)$:

$$\mu_\ell \omega_\ell^{(0)} - q_\ell \sum_{j=1}^r \mu_j \omega_j^{(0)} = 0,$$

thence, with

$$\Omega^{(0)}(\underline{y}, t) = \sum_{j=1}^r \mu_j \omega_j^{(0)}, \quad (3.17)$$

we obtain

$$\omega_\ell^{(0)} = \frac{q_\ell}{\mu_\ell} \Omega^{(0)}(\underline{y}, t). \quad (3.18)$$

In the summed equation (3.15) we have

$$\sum_{j=1}^r y_j \left[\mu_j \omega_j^{(0)} + \left(\frac{d\beta_j}{dt} - \lambda_j (\alpha_j - \beta_j) \right) \left(\sum_{\ell=1}^r \omega_\ell^{(0)} \right) \right] = 0. \quad (3.19)$$

Put

$$\varphi \equiv \varphi(\underline{\beta}, \underline{\alpha}) = \sum_{\ell=1}^r q_\ell / \mu_\ell; \quad (3.20)$$

then (3.19) becomes

$$\sum_{j=1}^r y_j \left[q_j + \left(\frac{d\beta_j}{dt} - \lambda_j (\alpha_j - \beta_j) \right) \varphi \right] \Omega^{(0)} = 0,$$

which is satisfied if

$$\begin{aligned} \frac{d\beta_j}{dt} &= \lambda_j (\alpha_j - \beta_j) - q_j / \varphi \\ &= \lambda_j (\alpha_j - \beta_j) - \mu_j \left(\frac{q_j / \mu_j}{\sum_{j=1}^r q_j / \mu_j} \right). \end{aligned} \quad (3.21)$$

The solution of (3.21) provides the initial approximation to the mean length of the j^{th} queue: $E[N_j(t)] \sim a\beta_j(t)$.

Note that $\chi^{(0)}(\underline{y}, t) = \sum_{\ell=1}^r \omega_\ell^{(0)} = \varphi \Omega^{(0)}(\underline{y}, t)$ provides the leading term in the asymptotic expansion for the ch.f. of $\underline{V}(t)$, the stochastic noise term that perturbs the mean and that $\chi^{(0)}(\underline{0}, t) = 1$.

Now return to (3.13) and isolate terms of order $1/\sqrt{a}$ to find that

$$\begin{aligned} \mu_\ell \omega_\ell^{(1)} - q_\ell \sum_{j=1}^r \mu_j \omega_j^{(1)} &= -i \sum_{j=1}^r \mu_j \left[q_\ell \omega_j^{(0)} y_j + \sum_{k=1}^r \frac{\partial q_\ell}{\partial \beta_k} \cdot \frac{\partial \omega_j^{(0)}}{\partial y_k} \right] \\ &\quad - i \sum_{j=1}^r \left[\frac{d\beta_j}{dt} - \lambda_j (\alpha_j - \beta_j) \right] y_j \omega_\ell^{(0)}; \end{aligned} \quad (3.22)$$

replace $\omega_j^{(0)}$ by $(q_j / \mu_j) \Omega^{(0)}$ and utilize $\sum_{j=1}^r q_j = 1$ and (3.21) to find

$$\mu_\ell \omega_\ell^{(1)} - q_\ell \sum_{j=1}^r \mu_j \omega_j^{(1)} = -i \left[\sum_{k=1}^r \frac{\partial q_\ell}{\partial \beta_k} \frac{\partial \Omega^{(0)}}{\partial y_k} + q_\ell \Omega^{(0)} \sum_{j=1}^r q_j y_j \right] + i \frac{q_\ell}{\mu_\ell} \frac{\Omega^{(0)}}{\varphi} \sum_{j=1}^r q_j y_j. \quad (3.23)$$

In order to make use of the summed equation, (3.15), first put $\Omega^{(1)}(\underline{y}, t) = \sum_{j=1}^r \mu_j \omega_j^{(1)}(\underline{y}, t)$ in (3.22), divide through by μ_ℓ and sum to obtain

$$\sum_{\ell=1}^r \omega_\ell^{(1)} = \varphi \Omega^{(1)} - i \left[\sum_{k=1}^r \frac{\partial \varphi}{\partial \beta_k} \frac{\partial \Omega^{(0)}}{\partial y_k} + \left(\varphi - \frac{R}{\varphi} \right) \Omega^{(0)} \sum_{k=1}^r q_k y_k \right] \quad (3.24)$$

in which

$$R = \sum_{\ell=1}^r q_\ell / \mu_\ell^2. \quad (3.25)$$

Multiply (3.23) by y_ℓ and sum:

$$\begin{aligned} \sum_{\ell=1}^r y_\ell \mu_\ell \omega_\ell^{(1)} &= \sum_{\ell=1}^r q_\ell y_\ell \sum_{j=1}^r \mu_j \omega_j^{(1)} - i \left[\sum_{\ell=1}^r \sum_{k=1}^r y_\ell \frac{\partial q_\ell}{\partial \beta_k} \frac{\partial \Omega^{(0)}}{\partial y_k} + \Omega^{(0)} \left(\sum_{j=1}^r q_j y_j \right)^2 \right] \\ &\quad + i \frac{\Omega^{(0)}}{\varphi} \sum_{\ell=1}^r \frac{q_\ell y_\ell}{\mu_\ell} \sum_{k=1}^r q_k y_k. \end{aligned} \quad (3.26)$$

Now utilize (3.26) and (3.24) in the summed equation, (3.15), and put

$$\Omega^{(1)} = \sum_{j=1}^r \mu_j \omega_j^{(1)} \quad (3.27)$$

to obtain

$$\begin{aligned}
& i \left\{ \frac{\Omega^{(1)} \sum_{\ell=1}^r q_{\ell} y_{\ell}}{\varphi} - i \left[\sum_{\ell=1}^r \sum_{k=1}^r y_{\ell} \frac{\partial q_{\ell}}{\partial \beta_k} \frac{\partial \Omega^{(0)}}{\partial y_k} + \Omega^{(0)} \left(\sum_{j=1}^r q_j y_j \right)^2 - \frac{\Omega^{(0)}}{\varphi} \sum_{\ell=1}^r \frac{q_{\ell} y_{\ell}}{\mu_{\ell}} \sum_{j=1}^r q_j y_j \right] \right. \\
& \quad \left. - \sum_{j=1}^r \frac{q_j y_j}{\varphi} \left(\frac{\varphi \Omega^{(1)}}{\varphi} - i \left[\sum_{k=1}^r \frac{\partial \varphi}{\partial \beta_k} \frac{\partial \Omega^{(0)}}{\partial y_k} + \varphi \Omega^{(0)} \sum_{k=1}^r q_k y_k - \frac{R}{\varphi} \Omega^{(0)} \sum_{k=1}^r q_k y_k \right] \right) \right\} \\
& \quad + \left\{ \sum_{j=1}^r \mu_j \omega_j^{(0)} \frac{y_j^2}{\varphi} + \sum_{j=1}^r \lambda_j (\alpha_j - \beta_j) \frac{y_j^2}{2} \left(\sum_{\ell=1}^r \omega_{\ell}^{(0)} \right) + \sum_{j=1}^r \lambda_j y_j \left(\sum_{\ell=1}^r \frac{\partial \omega_{\ell}^{(0)}}{\partial y_j} \right) + \sum_{\ell=1}^r \frac{\partial \omega_{\ell}^{(0)}}{\partial t} \right\} \\
& \quad = 0.
\end{aligned} \tag{3.28}$$

The terms designated by $\underline{\quad}$ and by $\underline{\underline{\quad}}$ cancel, leaving an equation involving only the first terms in the expansion (3.16). Further, in view of (3.18), the latter can be reduced to an equation involving only $\Omega^{(0)}(\underline{y}, t)$.

If we define

$$s_j(\underline{\beta}, \underline{\alpha}) = q_j(\underline{\beta}, \underline{\alpha}) / \varphi(\underline{\beta}, \underline{\alpha}), \quad r(\underline{\beta}, \underline{\alpha}) = R(\underline{\beta}, \underline{\alpha}) / \varphi(\underline{\beta}, \underline{\alpha}) \tag{3.29}$$

and recall, $\chi^{(0)}(\underline{y}, t) = \varphi(\underline{\beta}, \underline{\alpha}) \Omega^{(0)}(\underline{y}, t)$ then (3.28) is expressed as follows:

$$\begin{aligned}
\frac{\partial \chi^{(0)}}{\partial t} = & - \left(\sum_{j=1}^r \lambda_j y_j \frac{\partial \chi^{(0)}}{\partial y_j} + \sum_{j=1}^r \sum_{k=1}^r \frac{\partial s_j}{\partial \beta_k} y_j \frac{\partial \chi^{(0)}}{\partial y_k} \right) \\
& \left(-\frac{1}{2} \sum_{j=1}^r [\lambda_j (\alpha_j - \beta_j) + s_j] y_j^2 + \sum_{j=1}^r \sum_{k=1}^r \left[\frac{1}{\mu_j} - r(\underline{\beta}, \underline{\alpha}) \right] s_j s_k y_j y_k \right) \chi^{(0)}.
\end{aligned} \tag{3.30}$$

Equation (3.30) is recognizable as the partial differential equation satisfied by the ch.f. of the Ornstein-Uhlenbeck process. This shows that the scaled noise terms have normal/Gaussian marginal distributions, and permits the derivation of a system of differential equations for the covariance function of

the stochastic queue lengths. These are obtained by differentiating (3.30) at $\underline{y} = 0$: if $\rho_{j\ell}(t)$ is the limiting covariance of $V_j(t)$, $V_\ell(t)$, as defined in (3.3), then

$$\begin{aligned} & \frac{d}{dt} \rho_{j\ell}(t) + (\lambda_j + \lambda_\ell) \rho_{j\ell}(t) + \sum_{k=1}^r \left[\frac{\partial s_j}{\partial \beta_k} \cdot \rho_{\ell k}(t) + \frac{\partial s_\ell}{\partial \beta_k} \cdot \rho_{jk}(t) \right] \\ &= \left[2r(\underline{\beta}, \underline{\alpha}) - \frac{1}{\mu_j} - \frac{1}{\mu_\ell} \right] s_j s_\ell + [\lambda_j(\alpha_j - \beta_j) + s_j] \delta_{j\ell} \end{aligned} \quad (3.31)$$

with $\delta_{j\ell} = 1$ if $j = \ell$, $= 0$ if $j \neq \ell$ (Kronecker delta). We have used the fact that $\chi^{(0)}(\underline{0}, t) = 1$. A first-order correction term to the mean can also be derived, as can further terms in the asymptotic expansion: if

$$E[N_j(t)] = a\beta_j(t) + \sqrt{a}\xi_j(t) + o(1)$$

then

$$\frac{d\xi_j(t)}{dt} + \lambda_j \xi_j(t) + \sum_{k=1}^r \frac{\partial s_j}{\partial \beta_k} \xi_k(t) = 0; \quad (3.32)$$

note that if $\xi_j(0) = 0$, $j = 1, 2, \dots, r$, then $\xi_j(t) \equiv 0$.

It seems likely that under the normalization chosen the process $\underline{V}(t)$ actually converges weakly to the Ornstein-Uhlenbeck process, but we do not prove this fact here; see McNeil and Schach (1973) for an early approach to this issue.

We now turn to a numerical assessment of the quality of the approximation for the time-dependent mean and standard deviation of individual queue lengths, and a discussion of the effect of the several service scheduling rules.

4. NUMERICAL RESULTS

The asymptotic results will now be illustrated numerically, and compared to simulation. Two sample systems will be examined.

EXAMPLE 1: CONFIGURATION AND RATES

	Machines				
$j:$	1	2	3	4	5
$K_j:$	100	110	120	130	140
$\lambda_j:$	0.011	0.012	0.013	0.014	0.015
$v_j:$	3.0	3.0	3.0	3.0	3.0

The following abbreviations identify the various service disciplines and computational procedures in the tables:

- FCFS,A/S: First Come, First-Served; Analytical Method/Simulation
- LOLIN,A/S: Longest Line Next; Analytical Method/Simulation
- LAIN,A/S: Least Available Item Next; Analytical Method/Simulation

The simulation has been carried out so as to faithfully represent the service disciplines: FCFS,S literally keeps track of the order of arrival and serves accordingly, and LOLIN,S and LAIN,S likewise perform as stated when new service opportunities occur; they occur deterministically. The analytical procedures utilize the probabilistic schemes described. In all present cases initial conditions are nominally zero; in order to avoid problems with indeterminacy of q_i small values of $\beta_j(0)$ were utilized; weights w_j were taken to be equal. The three times selected to report results essentially cover the interesting range of temporal variation; the system is nearly in statistical equilibrium at the final value. Simulations were replicated 500 times; for details see Pilnick (1989). The typical simulation exercise for a single case, e.g.,

FCFS,S for all machines required about ten minutes of IBM 3033/4381 time, while the corresponding analytical-numerical solution was carried out in the order of a few seconds.

Table 1 allows comparison of the mean and standard deviation (parentheses) of queue lengths as (a) these measures are computed analytically-numerically, solving equations (3.20) and (3.26), and (b) by point-event simulation. It also invites comparison of the effect of the various disciplines.

Discussion

Figures in Table 1 show that

- For all disciplines employed, and at all times recorded, the analytical-numerical and simulated means and standard deviations agreed well. Of course the analytical-numerical computations were performed in a small fraction of simulation time. Note that the target p -value (tuning parameter) $p = 30$ provides numbers closer to simulation despite the danger of instability associated with very large p 's. Choice of a and of p must be guided by experiment at present.
- The stochastic variability around the time-dependent mean queue length, measured by the standard deviations, is always a small fraction of the mean. The FCFS standard deviations are always somewhat greater than those for the LOLIN or LAIN policies; these latter have a natural tendency to equalize queue lengths, both with respect to means *and* stochastic fluctuations. Studies not reported here, Pilnick (1989), suggest that the scaled marginal distribution of queue length, $V_j(t)$, is close to being normally distributed, as would be implied by the Ornstein-Uhlenbeck nature of the limiting ch.f., (3.30).
- The LAIN policy quickly adapts to equalize the number of unqueued, or available or idle machines in each group. This feature is of special interest when one available machine of each type is needed on a "platform" (e.g., ship, aircraft, *etc.*) in order that it be totally mission-ready. For other models and studies in this direction see Gaver *et al.*, (1991). Note that LAIN is myopic and makes no use of estimates of parameter values λ_j or μ_j , or even the number, r , of

competing machine types. The same general objective should be of interest if the machine types are components of a sequential manufacturing process and it is desired to avoid bottlenecks.

TABLE 1

MACHINE	SERVICE DISCIPLINES	TIMES		
		100	200	500
1	FCFS,S	40.5(5.3)	52.4(5.3)	57.2(5.1)
	FCFS,A	40.3(5.2)	52.6(5.3)	57.5(5.2)
	LOLIN,S	53.5(3.3)	68.7(3.1)	74.0(3.2)
	LOLIN,A	(p = 10) 50.9(3.7)	65.3(3.5)	70.8(3.4)
		(p = 30) 53.2(3.7)	68.0(3.4)	73.5(3.2)
	LAIN,S	NA	NA	NA
	LAIN,A	(p = 30) 33.1(3.7)	47.9(3.4)	53.5(3.2)
2	FCFS,S	47.4(5.5)	60.6(5.7)	65.5(5.6)
	FCFS,A	47.0(5.6)	60.5(5.6)	65.6(5.4)
	LOLIN,S	54.3(3.4)	69.2(3.1)	74.5(3.3)
	LOLIN,A	(p = 10) 53.0(3.9)	67.6(3.7)	73.1(3.6)
		(p = 30) 54.1(3.6)	68.6(3.4)	74.5(3.3)
	LAIN,S	NA	NA	NA
	LAIN,A	(p=30) 43.4(3.7)	58.1(3.4)	63.6(3.2)
3	FCFS,S	53.8(6.3)	68.5(5.8)	73.7(6.0)
	FCFS,A	54.0(5.9)	68.6(5.8)	73.8(5.7)
	LOLIN,S	54.8(3.5)	69.6(3.1)	74.9(3.1)
	LOLIN,A	(p = 10) 54.7(4.1)	69.6(3.9)	75.2(3.8)
		(p = 30) 53.7(3.7)	69.7(3.6)	75.3(3.5)
	LAIN,S	NA	NA	NA
	LAIN,A	(p = 30) 53.7(3.7)	68.2(3.4)	73.7(3.2)
4	FCFS,S	61.2(6.6)	76.9(6.1)	81.6(5.9)
	FCFS,A	61.2(6.2)	76.9(6.0)	82.3(5.8)
	LOLIN,S	55.4(3.6)	70.1(3.2)	75.3(3.3)
	LOLIN,A	(p = 10) 56.2(4.2)	71.3(4.1)	77.0(4.0)
		(p = 30) 55.2(3.7)	70.2(3.6)	75.9(3.5)
	LAIN,S	NA	NA	NA
	LAIN,A	(p = 30) 63.9(3.7)	78.4(3.4)	83.8(3.2)
5	FCFS,S	68.4(6.8)	84.4(6.1)	90.3(6.0)
	FCFS,A	68.8(6.5)	85.3(6.2)	90.8(6.0)
	LOLIN,S	55.9(3.6)	70.6(3.2)	75.8(3.3)
	LOLIN,A	(p = 10) 57.5(4.4)	72.9(4.3)	78.7(4.2)
		(p = 30) 55.7(3.8)	70.8(3.7)	76.5(3.5)
	LAIN,S	NA	NA	NA
	LAIN,A	(p = 30) 74.2(3.7)	88.5(3.4)	93.9(3.2)

EXAMPLE 2: CONFIGURATION AND RATES
(Failure and Repair Rates Differ)

	Machines				
j :	1	2	3	4	5
K_j :	100	110	120	130	140
λ_j :	0.015	0.020	0.025	0.030	0.035
ν_j :	1.0	1.1	1.2	1.3	1.4

Note that in this example the repair rates differ, whereas in the previous example they were the same. In Table 2 there appear the analytical and simulated means and standard deviation (parentheses) of the individual machine availabilities when the FCFS and LAIN service policies are in effect.

TABLE 2. MACHINES AVAILABLE

MACHINES	SERVICE DISCIPLINES	TIMES		
		50	100	200
1	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)
	LAIN,S	51.3(4.7)	24.5(3.9)	11.7(1.6)
	LAIN,A ($p = 10$)	52.1(4.8)	25.2(3.9)	11.5(1.7)
	FCFS,S	46.1(5.0)	22.8(4.5)	11.6(3.3)
	FCFS,A	46.5(5.2)	23.6(4.3)	12.5(3.3)
2	LAIN,S	43.9(4.3)	20.0(2.2)	11.3(1.6)
	LAIN,A ($p = 10$)	44.9(4.4)	20.2(2.6)	11.1(1.7)
	FCFS,S	41.0(5.2)	18.6(4.2)	10.7(3.3)
	FCFS,A	41.4(5.3)	18.9(4.0)	10.5(3.1)
	LAIN,S	40.1(3.4)	19.3(2.5)	11.0(1.7)
	LAIN,A ($p = 10$)	40.2(3.9)	19.2(2.5)	10.8(1.7)
3	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)
	LAIN,S	51.3(4.7)	24.5(3.9)	11.7(1.6)
	LAIN,A ($p = 10$)	52.1(4.8)	25.2(3.9)	11.5(1.7)
	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)
4	LAIN,S	51.3(4.7)	24.5(3.9)	11.7(1.6)
	LAIN,A ($p = 10$)	52.1(4.8)	25.2(3.9)	11.5(1.7)
	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)
	LAIN,S	51.3(4.7)	24.5(3.9)	11.7(1.6)
	LAIN,A ($p = 10$)	52.1(4.8)	25.2(3.9)	11.5(1.7)
5	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)
	LAIN,S	51.3(4.7)	24.5(3.9)	11.7(1.6)
	LAIN,A ($p = 10$)	52.1(4.8)	25.2(3.9)	11.5(1.7)
	FCFS,S	51.5(5.0)	28.9(4.6)	13.9(3.3)
	FCFS,A	52.1(5.0)	30.3(4.6)	16.1(3.7)

Discussion

Table 2 indicates that the analytical-numerical and simulation-generated means and standard deviations again agree reasonably well, even though individual type failure rates and repair rates differ. The major discrepancies seem to appear in FCFS discipline comparisons. The qualitative effects noted earlier are once again evident: generally small standard deviation to mean ratio (coefficient of variation); larger standard deviation to mean ratio (coefficient of variation) and larger standard deviation for FCFS than for LAIN, and a pronounced (an anticipated) tendency for LAIN to equalize the number of machines available as time advances.

Conclusions

Our conclusion is that asymptotic expressions based on the scaling (3.3) provide quite adequate approximations to the time-dependent or transient means and variances for the particular multitype repairman problems studied. These problems include attention to a variety of dynamic priority or service-adaptive policies via the device of tailored probabilistic selection of new service incumbents. Not surprisingly, convergence to the marginal distribution of an Ornstein-Uhlenbeck process occurs; actual weak convergence has not yet been demonstrated, but is a likely bonus.

REFERENCES

Ludwig Arnold (1974). *Stochastic Differential Equations, Theory and Applications*. Wiley-Interscience. John Wiley and Sons, N.Y.

D. R. Cox and W. L. Smith (1961). *Queues*. Methuen Monograph. London.

W. Feller (1967). *An Introduction to Probability Theory and its Applications*. John Wiley and Sons, N.Y. (Third Edition).

D. P. Gaver, K. Isaacson, S. E. Pilnick and R. Silveira (1991). Assessing and improving many-platform availability during intensive demand periods. (paper in preparation).

D. P. Gaver and P. A. Jacobs (1986). Processor-shared time-sharing models in heavy traffic. *SIAM J. Comput.*, **15**, pp. 1085-1100.

D. P. Gaver and J. A. Morrison (1991). Heavy-traffic analysis of multi-type queueing under probabilistically load-preferential service order. *SIAM J. Appl. Math.* (to appear).

M. P. Jaiswal, C. L. Sharma, Karmeshu (1990). A queueing model with hysteresis effect: an application to a machine interference problem. Unpublished paper.

D. R. McNeil and S. Schach (1973). Central limit analogues for Markov population processes, *J. of the Royal Stat. Soc. (B)*, No. 1, pp. 1-23.

J. A. Morrison (1987). Conditioned response-time distribution for a large closed processor-sharing system in very heavy usage. *SIAM J. Appl. Math.*, **47**, pp. 1117-1129.

S. E. Pilnick (1989). *Combat Logistics Problems*. Ph.D. Dissertation, Dept. of Operations Research, Naval Postgraduate School, Monterey, CA.

INITIAL DISTRIBUTION LIST

1.	Library (Code 0142).....	2
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
2.	Defense Technical Information Center.....	2
	Cameron Station	
	Alexandria, VA 22314	
3.	Office of Research Administration (Code 81)	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
4.	Prof. Peter Purdue.....	1
	Code OR-Pd	
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
5.	Department of Operations Research (Code 55).....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
6.	Prof. Donald Gaver, Code OR-Gv.....	15
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
7.	Prof. Patricia Jacobs	15
	Code OR/Jc	
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
8.	Center for Naval Analyses.....	1
	4401 Ford Avenue	
	Alexandria, VA 22302-0268	
9.	Dr. David Brillinger.....	1
	Statistics Department	
	University of California	
	Berkeley, CA 94720	

10. Dr. R. Gnanadesikan1
 Bellcore
 435 South Street
 Morris Township NJ 07960
11. Prof. Bernard Harris1
 Dept. of Statistics
 University of Wisconsin
 610 Walnut Street
 Madison, WI 53706
12. Prof. I. R. Savage.....1
 Dept. of Statistics
 Yale University
 New Haven, CT 06520
13. Prof. W. R. Schucany.....1
 Dept. of Statistics
 Southern Methodist University
 Dallas, TX 75222
14. Prof. D. C. Siegmund.....1
 Dept. of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305
15. Prof. H. Solomon1
 Department of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305
16. Dr. Ed Wegman.....1
 George Mason University
 Fairfax, VA 22030
17. Dr. P. Welch1
 IBM Research Laboratory
 Yorktown Heights, NY 10598
18. Dr. Neil Gerr1
 Office of Naval Research
 Arlington, VA 22217

19. Prof. Roy Welsch.....1
Sloan School
M.I.T.
Cambridge, MA 02139
20. Dr. J. Abrahams1
Code 1111, Room 607
Mathematical Sciences Division
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000
21. Prof. J. R. Thompson.....1
Dept. of Mathematical Science
Rice University
Houston, TX 77001
22. Dr. P. Heidelberger.....1
IBM Research Laboratory
Yorktown Heights
New York, NY 10598
23. Prof. M. Leadbetter.....1
Department of Statistics
University of North Carolina
Chapel Hill, NC 27514
24. Prof. D. L. Iglehart1
Dept. of Operations Research
Stanford University
Stanford, CA 94350
25. Prof. J. B. Kadane.....1
Dept. of Statistics
Carnegie-Mellon University
Pittsburgh, PA 15213
26. Prof. J. Lehoczky1
Department of Statistics
Carnegie-Mellon University
Pittsburgh, PA 15213

27. Dr. J. Maar (R513)1
National Security Agency
Fort Meade, MD 20755-6000
28. Prof. M. Mazumdar1
Dept. of Industrial Engineering
University of Pittsburgh
Pittsburgh, PA 15235
29. Prof. M. Rosenblatt1
Department of Mathematics
University of California, San Diego
La Jolla, CA 92093
30. Prof. H. Ciernoff1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138
31. Dr. T. J. Ott1
Bellcore
435 South Street
Morris Township, NJ 07960
32. Dr. Alan Weiss1
AT&T Bell Laboratories
Mountain Avenue
Murray Hill, NJ 07974
33. Prof. Joseph R. Gani1
Mathematics Department
University of California
Santa Barbara, CA 93106
34. Prof. Frank Samaniego1
Statistics Department
University of California
Davis, CA 95616
35. Dr. James McKenna1
AT&T Bell Laboratories
Mountain Avenue
Murray Hill, NJ 07974

36. Prof. R. Douglas Martin.....1
Department of Statistics, GN-22
University of Washington
Seattle, WA 98195

37. Dr. D. C. Hoaglin1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138

38. Prof. N. D. Singpurwalla1
George Washington University
Washington, DC 20052

39. Prof. George S. Fishman.....1
Curr. in OR & Systems Analysis
University of North Carolina
Chapel Hill, NC 20742

40. Dr. Alan F. Petty1
Code 7930
Naval Research Laboratory
Washington, DC 20375

41. Prof. Bradley Efron.....1
Statistics Dept.
Sequoia Hall
Stanford University
Stanford, CA 94305

42. Prof. Carl N. Morris.....1
Statistics Department
Harvard University
1 Oxford St.
Cambridge, MA 02138

43. Dr. John E. Rolph.....1
RAND Corporation
1700 Main St.
Santa Monica, CA 90406

44. Prof. Linda V. Green.....1
Graduate School of Business
Columbia University
New York, NY 10027

45. Dr. David Burman.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974

46. Dr. Ed Coffman.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974

47. Prof. William Jewell.....1
Operations Research Department
University of California, Berkeley
Berkeley, CA 94720

48. Prof. D. C. Siegmund.....1
Dept. of Statistics
Sequoia Hall
Stanford University
Stanford, CA 94305

49. Operations Research Center, Rm E40-164.....1
Massachusetts Institute of Technology
Attn: R. C. Larson and J. F. Shapiro
Cambridge, MA 02139

50. Arthur P. Hurter, Jr.1
Professor and Chairman
Dept. of Industrial Engineering and Management Sciences
Northwestern University
Evanston, IL 60201-9990

51. Institute for Defense Analysis.....1
1800 North Beauregard
Alexandria, VA 22311

52. Prof. J. W. Tukey1
Statistics Dept., Fine Hall
Princeton University
Princeton, NJ 08540
53. Dr. Daniel H. Wagner1
Station Square One
Paoli, PA 19301
54. Dr. Colin Mallows.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974
55. Dr. Jon Kettenring.....1
Bellcore
435 South Street
Morris Township, NJ 07960
56. Dr. S. R. Dalal.....1
AT&T Bell Telephone Laboratories
Mountain Avenue
Murray Hill, NJ 07974
57. Dr. M. J. Fischer1
Defense Communications Agency
1860 Wiehle Avenue
Reston, VA 22070
58. Dr. Prabha Kumar
Defense Communications Agency.....1
1860 Wiehle Avenue
Reston, VA 22070
59. Dr. B. Doshi1
AT&T Bell Laboratories
HO 3M-335
Holmdel, NJ 07733
60. Dr. D. M. Lucantoni.....1
AT&T Bell Laboratories
Holdmel, NJ 07733

61. Dr. V. Ramaswami1
MRE 2Q-358
Bell Communications Research, Inc.
435 South Street
Morristown, NJ 07960

62. Prof. G. Shantikumar1
The Management Science Group
School of Business Administration
University of California
Berkeley, CA 94720

63. Dr. D. F. Daley1
Statistic Dept. (I.A.S.)
Australian National University
Canberra, A.C.T. 2606
AUSTRALIA

64. Dr. Guy Fayolle1
I.N.R.I.A.
Dom de Voluceau-Rocquencourt
78150 Le Chesnay Cedex
FRANCE

65. Professor H. G. Daellenbach1
Department of Operations Research
University of Canterbury
Christchurch, NEW ZEALAND

66. Koh Peng Kong1
OA Branch, DSO
Ministry of Defense
Blk 29 Middlesex Road
SINGAPORE 1024

67. Professor Sir David Cox1
Nuffield College
Oxford, OX1 INF
ENGLAND

68. Dr. A. J. Lawrence.....1
 Department of Mathematical Statistics
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND

69. Dr. F. P. Kelly.....1
 Statistics Laboratory
 16 Mill Lane
 Cambridge
 ENGLAND


70. Dr. R. J. Gibbens.....1
 Statistics Laboratory
 16 Mill Lane
 Cambridge
 ENGLAND

71. Dr. John Copas.....1
 Dept. of Mathematical Statistics
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND

72. Dr. D. Vere-Jones.....1
 Dept. of Mathematics
 Victoria University of Wellington
 P. O. Box 196
 Wellington
 NEW ZEALAND

73. Prof. Guy Latouche.....1
 University Libre Bruxelles
 C. P. 212
 Blvd. De Triomphe
 B-1050 Bruxelles
 BELGIUM

DUDLEY KNOX LIBRARY



3 2768 00341860 9